

CANDIDATE  
NAME

--

CENTRE  
NUMBER

--	--	--	--	--

CANDIDATE  
NUMBER

--	--	--	--

**CHEMISTRY**

**5070/21**

Paper 2 Theory

**October/November 2015**

**1 hour 30 minutes**

Candidates answer on the Question Paper.

No Additional Materials are required.

**READ THESE INSTRUCTIONS FIRST**

Write your Centre number, candidate number and name on all the work you hand in.

Write in dark blue or black pen.

You may use an HB pencil for any diagrams or graphs.

Do not use staples, paper clips, glue or correction fluid.

**DO NOT WRITE IN ANY BARCODES.**

**Section A**

Answer **all** questions.

Write your answers in the spaces provided in the Question Paper.

**Section B**

Answer any **three** questions.

Write your answers in the spaces provided in the Question Paper.

Electronic calculators may be used.

You may lose marks if you do not show your working or if you do not use appropriate units.

A copy of the Periodic Table is printed on page 20.

At the end of the examination, fasten all your work securely together.

The number of marks is given in brackets [ ] at the end of each question or part question.

This document consists of **17** printed pages and **3** blank pages.

**Section A**

Answer **all** the questions in this section in the spaces provided.

The total mark for this section is 45.

**A1** Choose from the following gases to answer the questions below.

**ammonia**  
**argon**  
**carbon dioxide**  
**chlorine**  
**ethane**  
**ethene**  
**nitrogen**  
**nitrogen(II) oxide**  
**oxygen**  
**sulfur dioxide**

Each of these gases can be used once, more than once or not at all.

Which gas

**(a)** forms approximately 1% of the air,

.....[1]

**(b)** bleaches damp litmus paper,

.....[1]

**(c)** reacts with water to form an alkaline solution,

.....[1]

**(d)** can undergo polymerisation,

.....[1]

**(e)** is a compound formed in the atmosphere as a result of lightning activity,


.....[1]

**(f)** is a diatomic molecule containing a total of 16 electrons?

.....[1]

[Total: 6]

- A2** Coal is a mixture of carbon compounds with a small amount of sulfur.  
A sample of coal is heated in the absence of air using the apparatus shown.



The distillate is a mixture of a colourless liquid and tar.

- (a)** The colourless liquid contains ammonia,  $\text{NH}_3$ .

Draw a 'dot-and-cross' diagram for ammonia.  
Show only the outer shell electrons.

[2]

- (b)** The tar contains ethanoic acid.  
When warmed in the presence of sulfuric acid, ethanoic acid reacts with propanol to form an ester.

Name and draw the structure of this ester showing all the atoms and all the bonds.

name.....

structure

[2]

(c) The tar also contains a compound with the following composition.

element	percentage by mass
carbon	76.60
hydrogen	6.38
oxygen	17.02

Deduce the empirical formula of this compound.

empirical formula .....[2]

(d) When coal is burned, an acidic gas is produced which decolourises acidified aqueous potassium manganate(VII). This gas contributes to acid rain.

(i) Identify this gas and describe how acid rain is formed.

.....  
 .....  
 .....[2]

(ii) Give one adverse effect of acid rain on buildings.

.....[1]

(iii) Acid rain can have an adverse effect on respiration.

Write an equation to represent the process of respiration.

.....[2]

[Total: 11]

**A3** A layer of ozone is present in the stratosphere about 30 km above the Earth's surface.

**(a)** Compounds with formulae such as  $\text{CCl}_3\text{F}$  and  $\text{C}_2\text{ClF}_5$  are responsible for the depletion of ozone.

**(i)** Give the general name for these compounds.

.....[1]

**(ii)** Explain, in terms of human health, why it is important that the ozone in the stratosphere does not become too depleted.

.....

.....[2]

**(b)** In the stratosphere, ozone,  $\text{O}_3$ , is broken down to oxygen by photochemical reactions.

**(i)** What is meant by the term *photochemical reaction*?

.....


.....[1]

**(ii)** Construct an equation for the breakdown of ozone molecules to oxygen molecules.

.....[1]

**(c)** Ozone oxidises  $\text{Fe}^{2+}$  ions to  $\text{Fe}^{3+}$  ions.

Complete the ionic equation for this reaction.



[Total: 6]

**A4** Copper is a metal.

**(a)** Draw a labelled diagram to show the bonding in copper.

[2]

**(b)** Explain why metals are malleable.

.....  
.....  
.....[2]

**(c)** Copper corrodes slowly in damp air.

One of the corrosion products has the formula  $\text{CuCO}_3 \cdot \text{Cu}(\text{OH})_2$ .



**(i)** Calculate the percentage by mass of copper in this compound.

[2]

**(ii)** How could you show that  $\text{CuCO}_3 \cdot \text{Cu}(\text{OH})_2$  contains carbonate ions?

.....  
.....  
.....[2]

- (d) Copper is oxidised by concentrated sulfuric acid.  
This redox reaction can be represented by equations **A** and **B**.



Which reaction, **A** or **B**, is oxidation and which is reduction? Explain your answer.


.....  
.....  
.....[2]

[Total: 10]





**A6** When 1 mole of sodium hydroxide reacts with excess hydrochloric acid, 57.1 kJ of energy is released.



(a) Calculate the energy released when 12.0g of sodium hydroxide reacts with excess hydrochloric acid.

[2]

(b) Calculate the volume of 0.200 mol/dm<sup>3</sup> HCl which contains 2.19g of HCl.

[2]

(c) Aqueous hydrochloric acid contains chloride ions.

Describe a test for chloride ions.

test .....

result ..... [2]

(d) Zinc oxide reacts with both hydrochloric acid and sodium hydroxide.

Which term describes this behaviour of zinc oxide?

..... [1]


[Total: 7]

**Section B**

Answer **three** questions from this section in the spaces provided.

The total mark for this section is 30.

**B7** The structures of graphite and boron nitride are shown below.



**(a)** Like graphite, boron nitride feels slippery to the touch.

Explain, in terms of structure and bonding, why boron nitride feels slippery to the touch.

.....

.....

.....[2]

**(b)** An isotope of boron is represented by the symbol  $^{11}_5\text{B}$ .

Deduce the number of protons and neutrons in this isotope of boron.

number of protons .....

number of neutrons .....[1]


**(c)** Explain why graphite has a high melting point.

.....

.....

.....[2]

(d) Dilute sulfuric acid can be electrolysed using graphite electrodes.



(i) Graphite is a good electrical conductor.

Explain why graphite conducts electricity.

.....[1]

(ii) Give another property of graphite that makes it useful as an electrode in this electrolysis.

.....[1]

(e) During the electrolysis of dilute sulfuric acid, oxygen is released at the anode (positive electrode) and hydrogen is released at the cathode (negative electrode).

(i) Complete the equation for the reaction at the anode.




(ii) Construct the equation for the reaction at the cathode.

.....[1]

(iii) Explain why the volume of hydrogen produced is approximately double that of the oxygen.

.....  
 .....[1]

**B8** A sample of 0.030g of small pieces of magnesium is added to 20cm<sup>3</sup> of 0.10mol/dm<sup>3</sup> hydrochloric acid.




**(a) (i)** Show by calculation which reactant is in excess.

[3]

**(ii)** What would you observe in this reaction?

.....[1]

- (b) The graph shows how the volume of hydrogen gas produced, at room temperature and pressure, changes with time.



Calculate the total mass of hydrogen formed.

mass of hydrogen = ..... g [2]

- (c) The experiment is repeated at the same temperature and pressure. The same mass of magnesium is added but magnesium powder is used instead of small pieces.

The rate of reaction is faster. Explain why.

.....  
 .....  
 ..... [2]

- (d) Magnesium reacts with nitrogen when heated. The ionic compound magnesium nitride,  $Mg_3N_2$ , is formed.

(i) Construct the equation, including state symbols, for this reaction.


..... [1]

(ii) Deduce the charge on the nitride ion.

..... [1]

[Total: 10]

**B9** The structure of glycolic acid can be represented as shown.



**(a)** Glycolic acid is a solid at room temperature.

Describe the arrangement and motion of the molecules in glycolic acid at room temperature.

arrangement .....

motion ..... [2]

**(b)** Glycolic acid can polymerise with itself to form a polyester called poly(glycolic acid).


**(i)** What type of polymer is a polyester?

..... [1]

**(ii)** Draw a section of the polymer chain of poly(glycolic acid) showing at least two repeating units.

[2]

- (c) Glycolic acid is produced by heating methanal, carbon monoxide and water in the presence of a sulfuric acid catalyst.



- (i) A sample of 1800g of methanal reacts with excess carbon monoxide and water. The percentage yield of glycolic acid is 45%.

Calculate the mass, in grams, of glycolic acid produced.

mass of glycolic acid = ..... g [3]


- (ii) Glycolic acid is a weak acid. Sulfuric acid is a strong acid.

Explain the difference between a strong acid and a weak acid.

.....  
 .....  
 ..... [2]

[Total: 10]

**B10** Methanol is manufactured by reacting carbon monoxide with hydrogen. The forward reaction is exothermic.



- (a) Predict and explain the effect of increasing the pressure on the position of equilibrium. The temperature remains constant.

.....  
 .....  
 ..... [2]

- (b) Predict and explain the effect of decreasing the temperature on the position of equilibrium. The pressure remains constant.

.....  
 .....  
 ..... [2]

- (c) The rate of reaction decreases when the temperature is lowered.

Explain why.

.....  
 .....  
 ..... [2]

- (d) The reaction is catalysed by copper.

- (i) Describe and explain the effect of a catalyst on this reaction.

.....  
 .....  
 ..... [2]

- (ii) Copper is a transition element. Many transition elements are catalysts.

Give **two** other properties of copper that identify it as a transition element.

.....  
 ..... [2]

[Total: 10]







**BLANK PAGE**

---

Permission to reproduce items where third-party owned material protected by copyright is included has been sought and cleared where possible. Every reasonable effort has been made by the publisher (UCLES) to trace copyright holders, but if any items requiring clearance have unwittingly been included, the publisher will be pleased to make amends at the earliest possible opportunity.

To avoid the issue of disclosure of answer-related information to candidates, all copyright acknowledgements are reproduced online in the Cambridge International Examinations Copyright Acknowledgements Booklet. This is produced for each series of examinations and is freely available to download at [www.cie.org.uk](http://www.cie.org.uk) after the live examination series.

Cambridge International Examinations is part of the Cambridge Assessment Group. Cambridge Assessment is the brand name of University of Cambridge Local Examinations Syndicate (UCLES), which is itself a department of the University of Cambridge.

**DATA SHEET**  
**The Periodic Table of the Elements**

		Group																																			
		I	II	III	IV	V	VI	VII	0																												
		<table border="1" style="width: 100%; border-collapse: collapse; text-align: center;"> <tr> <td style="width: 10%;">1</td> <td style="width: 10%;"><b>H</b> Hydrogen 1</td> <td colspan="8"></td> <td style="width: 10%;">4</td> <td style="width: 10%;"><b>He</b> Helium 2</td> </tr> </table>										1	<b>H</b> Hydrogen 1									4	<b>He</b> Helium 2														
1	<b>H</b> Hydrogen 1									4	<b>He</b> Helium 2																										
7	<b>Li</b> Lithium 3	9	<b>Be</b> Beryllium 4	11	<b>B</b> Boron 5	12	<b>C</b> Carbon 6	13	<b>Al</b> Aluminium 13	14	<b>N</b> Nitrogen 7	15	<b>O</b> Oxygen 8	16	<b>F</b> Fluorine 9	17	<b>Ne</b> Neon 10																				
23	<b>Na</b> Sodium 11	24	<b>Mg</b> Magnesium 12	25	<b>Mn</b> Manganese 25	26	<b>Fe</b> Iron 26	27	<b>Co</b> Cobalt 27	28	<b>Ni</b> Nickel 28	29	<b>Cu</b> Copper 29	30	<b>Zn</b> Zinc 30	31	<b>Ga</b> Gallium 31	32	<b>Ge</b> Germanium 32	33	<b>As</b> Arsenic 33	34	<b>Se</b> Selenium 34	35	<b>Br</b> Bromine 35	36	<b>Kr</b> Krypton 36										
39	<b>K</b> Potassium 19	40	<b>Ca</b> Calcium 20	41	<b>Nb</b> Niobium 41	42	<b>Mo</b> Molybdenum 42	43	<b>Tc</b> Technetium 43	44	<b>Ru</b> Ruthenium 44	45	<b>Rh</b> Rhodium 45	46	<b>Pd</b> Palladium 46	47	<b>Ag</b> Silver 47	48	<b>Cd</b> Cadmium 48	49	<b>In</b> Indium 49	50	<b>Sn</b> Tin 50	51	<b>Sb</b> Antimony 51	52	<b>Te</b> Tellurium 52	53	<b>I</b> Iodine 53	54	<b>Xe</b> Xenon 54						
85	<b>Rb</b> Rubidium 37	86	<b>Sr</b> Strontium 38	87	<b>Y</b> Yttrium 39	88	<b>Zr</b> Zirconium 40	89	<b>Hf</b> Hafnium 72	90	<b>Ra</b> Radium 88	91	<b>Ta</b> Tantalum 73	92	<b>W</b> Tungsten 74	93	<b>Re</b> Rhenium 75	94	<b>Os</b> Osmium 76	95	<b>Pt</b> Platinum 78	96	<b>Au</b> Gold 79	97	<b>Hg</b> Mercury 80	98	<b>Tl</b> Thallium 81	99	<b>Pb</b> Lead 82	100	<b>Bi</b> Bismuth 83	101	<b>Po</b> Polonium 84	102	<b>At</b> Astatine 85	103	<b>Rn</b> Radon 86
133	<b>Cs</b> Caesium 55	137	<b>Ba</b> Barium 56	138	<b>La</b> Lanthanum 57	139	<b>Ce</b> Cerium 58	140	<b>Pr</b> Praseodymium 59	141	<b>Nd</b> Neodymium 60	142	<b>Pm</b> Promethium 61	143	<b>Sm</b> Samarium 62	144	<b>Eu</b> Europium 63	145	<b>Gd</b> Gadolinium 64	146	<b>Tb</b> Terbium 65	147	<b>Dy</b> Dysprosium 66	148	<b>Ho</b> Holmium 67	149	<b>Er</b> Erbium 68	150	<b>Tm</b> Thulium 69	151	<b>Yb</b> Ytterbium 70	152	<b>Lu</b> Lutetium 71	153	<b>Yt</b> Ytterbium 71	154	<b>La</b> Lanthanum 71
223	<b>Fr</b> Francium 87	226	<b>Ra</b> Radium 88	227	<b>Ac</b> Actinium 89	228	<b>Th</b> Thorium 90	229	<b>Pa</b> Protactinium 91	230	<b>U</b> Uranium 92	231	<b>Np</b> Neptunium 93	232	<b>Pu</b> Plutonium 94	233	<b>Am</b> Americium 95	234	<b>Cm</b> Curium 96	235	<b>Bk</b> Berkelium 97	236	<b>Cf</b> Californium 98	237	<b>Es</b> Einsteinium 99	238	<b>Fm</b> Fermium 100	239	<b>Md</b> Mendelevium 101	240	<b>No</b> Nobelium 102	241	<b>Lr</b> Lawrencium 103	242	<b>Lu</b> Lutetium 103	243	<b>Lu</b> Lutetium 103

\* 58–71 Lanthanoid series  
† 90–103 Actinoid series

**Key**

a	<b>X</b>
b	†

a = relative atomic mass  
X = atomic symbol  
b = atomic (proton) number

The volume of one mole of any gas is 24 dm<sup>3</sup> at room temperature and pressure (r.t.p.).